

# Aerosol transmission of SARS-CoV-2: inhalation as well as exhalation matters

Authors: Keir EJ Philip<sup>1,4\*</sup>, Adam Lewis<sup>2</sup>, Sara C Buttery<sup>1,4</sup>, Colm McCabe<sup>1,4</sup>, Daisy Fancourt<sup>3</sup>, Christopher Orton<sup>1,4</sup>, Michael I Polkey<sup>1,4</sup>, Nicholas S Hopkinson<sup>1,4</sup>

\*Corresponding author

Email: k.philip@imperial.ac.uk

Address: Muscle Lab, Royal Brompton Hospital, London, SW3 6HP

Keir Philip ORCID: <https://orcid.org/0000-0001-9614-3580>

## Affiliations:

- 1) National Heart and Lung Institute, Imperial College London, London, United Kingdom
- 2) Department of Health Sciences, Brunel University, London, United Kingdom
- 3) Department of Behavioural Science and Health, University College London, London, United Kingdom
- 4) Royal Brompton and Harefield NHS Foundation Trust, London, United Kingdom

**Author contributions:** All others contributed to the manuscript concept and writing, and all agreed on the final version.

**Running head:** SARS-CoV-2: inhalation as well as exhalation matters

**Declaration of interests:** No competing interests.

**Funding:** Keir Philip is supported by the Imperial College Clinician Investigator Scholarship. The funder had no role in the writing of, or decision to submit, this manuscript.

**Word Count:** 448

We read with great interest the article by Etchernach and colleagues(1) on the topic of aerosol dispersion during singing and speaking as a potential COVID-19 transmission pathway. In the article, as has been the case more broadly regarding this mode of transmission, attention has focused on factors that influence emission of virus (i.e. aerosol production by the infected individual) when singing or speaking. However, the ventilatory pattern of individuals exposed to aerosolised virus is also an important factor as this is likely to be a key modulator of the 'dose' of virus-containing aerosol inhaled. As such, inclusion of such parameters in discussion regarding aerosol transmission is important when considering why certain contexts such as choirs, restaurants, and bars, where speaking, singing and shouting are common, have been linked to infection clusters(2). Such an appreciation may reframe the discussion to include 'super-receptiveness' as a component of 'super-spreader' events.

Ventilatory parameters vary greatly depending on both the type and intensity of activity and should feature more prominently when considering aerosol transmission. We recently investigated the physiological demands of 'Singing for Lung Health' (SLH) in healthy volunteers(3), and found that when participating in the singing component of the protocol, minute ventilation increased from resting volumes of 11L/min (9-13) (median, interquartile range (IQR)) to 22.42L/min (IQR 16.83 - 30.54); and the median volume per breath increased from 0.69L (IQR 0.63 - 0.77) to 2.11L (IQR 1.92 - 2.70). Other researchers, comparing talking with quiet breathing, found increases in parameters including minute ventilation, tidal volumes, and breathing frequency(4, 5).

Both increased minute ventilation and increased tidal volume are likely to be relevant to aerosol transmission. Firstly, the more aerosolised viral particles inhaled the larger the inoculum, which will impact the chance of developing disease, and may also influence disease severity(6). Secondly, greater inhalation will increase the alveolar area exposed to virus containing aerosols which may have implications for the viral processing and the immune response(7).

Considering patterns of inhalation as well as exhalation should enable a more complete appreciation of context specific viral transmission dynamics. This is particularly relevant to contexts where minute ventilation is increased due to physical activity (gyms supermarkets etc.), but also where groups of people are vocalising, such as choral singing, restaurants, bars, and sports crowds. Appreciating these factors does not change the fundamental focus of mitigation measures of hygiene, face coverings, physical distance, and avoiding contexts with poor ventilation. However, given these considerations, particularly with new more infectious SARS-CoV-2 variants in circulation, there may now be a stronger argument for face coverings that reduce risk of inhaling aerosols rather than just reducing their emission, especially in contexts where people are vocalising, exercising, or other risk factors are present.

## **References**

1. Echternach M, Gantner S, Peters G, Westphalen C, Benthous T, Jakubass B, Kuranova L, Dollinger M, Kniesburges S. Impulse Dispersion of Aerosols during Singing and Speaking: A Potential COVID-19 Transmission Pathway. *American journal of respiratory and critical care medicine* 2020; 202: 1584-1587.
2. Jones NR, Qureshi ZU, Temple RJ, Larwood JPJ, Greenhalgh T, Bourouiba L. Two metres or one: what is the evidence for physical distancing in covid-19? *Bmj* 2020; 370: m3223.
3. Philip KEJ, Lewis A, Buttery SC, McCabe C, Fancourt D, Orton C, Polkey MI, Hopkinson NS. The physiological demands of Singing for Lung Health compared to treadmill walking. *medRxiv* 2020.
4. Russell BA, Cerny FJ, Stathopoulos ET. Effects of varied vocal intensity on ventilation and energy expenditure in women and men. *J Speech Lang Hear Res* 1998; 41: 239-248.
5. Hoit JD, Lohmeier HL. Influence of continuous speaking on ventilation. *J Speech Lang Hear Res* 2000; 43: 1240-1251.
6. Guallar MP, Meirino R, Donat-Vargas C, Corral O, Jouve N, Soriano V. Inoculum at the time of SARS-CoV-2 exposure and risk of disease severity. *Int J Infect Dis* 2020; 97: 290-292.
7. Chu H, Chan JF, Wang Y, Yuen TT, Chai Y, Hou Y, Shuai H, Yang D, Hu B, Huang X, Zhang X, Cai JP, Zhou J, Yuan S, Kok KH, To KK, Chan IH, Zhang AJ, Sit KY, Au WK, Yuen KY. Comparative Replication and Immune Activation Profiles of SARS-CoV-2 and SARS-CoV in Human Lungs: An Ex Vivo Study With Implications for the Pathogenesis of COVID-19. *Clin Infect Dis* 2020; 71: 1400-1409.

**Acknowledgments:**

KP would like to acknowledge the National Institute for Health Research (NIHR) Biomedical Research Centre based at Imperial College Healthcare NHS Trust and Imperial College London for their support. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.